

NIPT REQUEST FORM

Cell-free DNA prenatal test

Mr. Colin Conaghan
FRCOG, FRANZCOG.

Dr. Angela Beard
FRANZCOG

Dr. Geeta Singh
FRANZCOG

CONTACT US

Hiatt Chambers
St George's Medical Centre
249 Papanui Rd
Christchurch

Phone 03 3754060
Fax 03 3754061
enquire@coga.org.nz

www.coa.org.nz



Patient Details

Name..... DOB.....

Address.....

.....

Contact Phone No..... NHI.....

Referrer Details

Doctor / Midwife..... Registration No.....

Address.....

.....

Contact Phone No..... Fax No.....

Signature..... Date.....

Clinical Details

LMP..... EDD..... Blood Group.....

Maternal age..... Height..... cm

NT scan abnormal or high risk MSS1..... Weight..... kg

Previous genetic abnormality

Maternal request for NIPT

Singleton Pregnancy

Twin Pregnancy

Notes

NIPT blood test does not replace 12 week NT scan.

Invasive testing (CVS/amniocentesis) recommended if:

1. Nuchal translucency >3.5mm
2. Fetal structural Lesion

Fetal echocardiography at a tertiary centre recommended if NT >3.5mm.

12 week scan will effectively screen for open spina bifida.

12 week scan will screen for some major structural anomalies.

Health Benefits (Section 88) claim available for
• NIPT specialist consultation and genetic advice

Percent Detection rate:

- trisomy 21 >99%
- trisomy 18 >98%
- trisomy 13 >98%
- sex chromosome anomalies:
depends on anomaly (Turner Synd >95%)
- gender >99%